

Cite this article as: Tsang W, Modi A, Ahmed I, Ohri SK. Do external support devices reduce sternal wound complications after cardiac surgery? *Interact CardioVasc Thorac Surg* 2016; doi:10.1093/icvts/iww270.

Do external support devices reduce sternal wound complications after cardiac surgery?

William Tsang^a, Amit Modi^{b,*}, Ishtiaq Ahmed^b and Sunil K. Ohri^c

^a The University of Manchester Medical School, Manchester, UK

^b Department of Cardiac Surgery, Royal Sussex County Hospital, Brighton, UK

^c Wessex Cardiothoracic Centre, Southampton University Hospitals, Southampton, UK

* Corresponding author. Department of Cardiac Surgery, Royal Sussex County Hospital, Brighton BN2 5BE, UK. Tel: +44-1273-696955; fax: +44-1273-664464; e-mail: drmodiamit@hotmail.com (A. Modi).

Received 26 December 2015; received in revised form 25 June 2016; accepted 13 July 2016

Abstract

A best evidence topic in cardiac surgery was written according to a structured protocol. The question addressed was whether external support devices reduce sternal wound complications after cardiac surgery with sternotomy. Altogether 116 papers were found using the reported search, of which six presented the best evidence to answer the clinical question. The author, year, journal, study type, patient group studied, relevant outcomes, results and study weaknesses are tabulated. Six randomized controlled trials investigating the effect of external chest support devices on sternal wound complications in adult patients undergoing sternotomy for cardiac surgery were selected. These studies demonstrate a significant reduction of deep sternal wound complication on comparing external support with no support. Non-elastic devices were more effective in reducing sternal complication compared with the elastic bandage (four trials). Three studies reported significant reduction of mean hospital stay in patients receiving non-elastic chest support devices. We conclude that early post-sternotomy use of an external non-elastic sternal support device reduces overall sternal wound complications and may reduce the hospital length of stay.

Keywords: Postoperative complication • Wound infection • Cardiac surgical procedures • External chest support • Review

INTRODUCTION

A best evidence topic was constructed according to a structured protocol. This protocol is fully described in the *ICVTS* [1].

CLINICAL SCENARIO

During the morning HDU round, the consultant surgeon suggests using an external thoracic support device on a male patient with moderate chronic obstructive pulmonary disease (COPD) who underwent a triple vessel coronary artery bypass surgery yesterday. The nurse accompanying you promptly applies the device but on the following day, you find the patient not wearing the device. On direct questioning, the patient says that the device is uncomfortable and would wear it only if there was sufficient evidence to support reduced sternal wound complication with its use. Therefore, you resolve to search the literature to find the evidence.

THREE-PART QUESTION

Do [external support devices] reduce [sternal wound complications] in patients undergoing [cardiac surgery with sternotomy]?

SEARCH STRATEGY

MEDLINE and EMBASE until April 2016 using the OVID interface. [exp. Cardiac surgical procedures/OR exp. thoracic surgery/OR exp. sternotomy/OR exp. sternum/OR thorax.mp.] AND [exp. dehiscence/OR exp. postoperative pain/OR infection.mp./OR instability.mp./OR mediastinitis.mp./OR wound.mp.] AND (corset.mp./OR brace.mp./OR harness.mp./OR bra.mp./OR vest.mp./OR external support.mp./OR thorax support.mp./OR chest support.mp./OR binder.mp./OR Stern-E-Fix.mp./OR Posthorax.mp./or elastic bandage.mp./or Cardibra.mp./or Sternshield.mp.)

SEARCH OUTCOME

The search returned 136 articles and after de-duplication 116 articles remained, of which 6 were included in the BET analysis reported below. The relevant papers are presented in Table 1.

COMMENTS

Six clinical studies, all randomized controlled trials (RCTs), involving 5826 adult patients were found suitable. These studies

Table 1: Best evidence papers

Author, year, journal, study type	Population group	Outcome	Key results	Comments
Gorlitzer et al. (2009), <i>Eur J Cardiothorac Surg</i> [2] Randomized controlled trial, single centre	455 patients, male and female gender, randomized immediately after cardiac surgery with sternotomy to receive Posthorax vest or no vest	All sternal wound complications	With vest = 0.6% Without vest = 4.9% A versus B (P = 0.015)	Sternal closure technique and perioperative use of antibiotics similar
	Group A: Vest group (n = 175)	Sternal wound dehiscence	Refused vest = 9.4% A versus C (P = 0.003)	Vest was applied within 24 h
	Group B: No Vest group (n = 227)	Superficial sternal complication	With vest = 0 Without vest = 0.4% Refused vest = 0	Diabetes mellitus was significantly lower in the 'No vest group'
	Group C: Refused after randomization (n = 53)	Deep sternal wound infection	With vest = 0.6% Without vest = 1.3% Refused vest = 1.9%	Method of randomization not mentioned
	Exclusion criteria: Less than 20 years old, congenital heart defects or mechanical reanimation or previous chest irradiation	Postoperative pain (visual analogue scale)	With vest = 0 Without vest = 1% Refused vest = 7.5%	23.2% of patients refused to wear the vest due to comfort issues
	Follow-up: 90 days	Length of hospital stay (days)	No difference (numerical data not provided)	
			With vest = 12 Without vest = 11 Refused vest = 12 (P = 0.53)	
Celik et al. (2011), <i>J Thorac Cardiovasc Surg</i> [3] This paper reports 2 studies. <i>Study 1</i> Retrospective cohort <i>Study 2</i> Randomized controlled trial, single centre	<i>Study 1</i> 842 patients who had undergone elective cardiac surgery with sternotomy	<i>Study 1</i> Sternal dehiscence and deep sternal wound infection	<i>Study 1</i> Significantly higher in COPD group (7.9 vs 1.2%; P < 0.001)	<i>Study 1</i> Figure-of-eight wire closure used in all patients
	Group 1a: COPD (n = 328)		COPD severity had significant effect (P = 0.002)	
	Group 1b: No COPD (n = 514)			
	<i>Study 2</i> 221 patients with moderate to severe COPD undergoing cardiac surgery with sternotomy	<i>Study 2</i> All sternal wound complications	<i>Study 2</i> With vest = 1% Without vest = 11.6% (P = 0.002)	<i>Study 2</i> Robiscek closure for all patients
	Group 2a: Vest group (n = 100)	Sternal wound dehiscence	With vest = 1% Without vest = 2.5% (P = 0.628)	Well-matched groups
	Group 2b: No vest group (n = 121)	Superficial sternal complication	With vest = 0 Without vest = 2.5% (P = 0.253)	Well powered for deep sternal wound complications
	Patients followed for 6 months postoperatively	Deep sternal wound infection	With vest = 0 Without vest = 6.6% (P = 0.009)	Highlights the impracticality to adjust the device for use in obese female patients
	Duration of hospital stay (days)	With vest = 13.7 Without vest = 17.8 (P = 0.03)		
	Number needed to treat	15 patients with moderate to severe COPD		
Naismith and Street (2005), <i>Eur J Cardiovasc Nurs</i> [4] Randomized controlled trial, single centre	20 female patients with bra cup size \geq C cup were randomly allocated into Cardibra or regular bra after cardiac surgery	Pain scores (Likert scale)	No significant difference	Extremely small sample size limiting statistical analysis
	Group 1: Cardibra (n = 10)	Sternal wound dehiscence or infection	None in either group	Cardibra applied immediately after surgery, regular bra applied 3 days postoperatively
	Group 2: Regular bra (n = 10)	Swelling	No swelling in treatment group Control group 27 mm ² at 7 days and 10.6 mm ² at 42 days	Use of analgesia not analysed
	Exclusion criteria: Previous mastectomy, pregnancy, emergency, unable to comply with follow-up protocol (6 weeks)	Comfort	No significant difference. (P-values not provided)	Comfort scores were primitive and could be more sophisticated

Continued

Table 1: (Continued)

Author, year, journal, study type	Population group	Outcome	Key results	Comments
Gorlitzer et al. (2010), <i>Interact CardioVasc Thorac Surg</i> [5] Randomized controlled trial, multicentre	1814 patients, male and female gender, randomized immediately after cardiac surgery with sternotomy to receive Posthorax vest or elastic chest bandage	Total complication rate: (Patients requiring additional sternal procedures)	Vest = 0.61% Bandage = 3.87% ($P = 0.047$)	Multicentre trial Cefazolin 1 g given IV for 48–72 h or until chest drain removed
	Group A: Chest elastic bandage ($n = 905$)	Sternal dehiscence	Vest = 0 Bandage = 0.77% ($P = 0.046$)	Vest applied after 48 h and patients advised to wear vest for 6 weeks
	Group B: Thorax vest 909 allocated 254 excluded ($n = 655$)	Superficial wound infections	Vest = 0.61% Bandage = 1.11% ($P = 0.42$)	Patients who failed to use vest were excluded (27.9%)
		Deep sternal infections	Vest = 0 Bandage = 1.99% ($P = 0.0001$)	Rate of AVR higher in elastic bandage group
		Hospitalization time (days)	Vest = 14.8 Bandage = 17.3 ($P = 0.04$)	
	1560 included in final analysis	ICU time (days)	Vest = 3.1 Bandage = 2.6 ($P = 0.12$)	
	No significant differences between demographic or risk factors of groups and STS infection risk score equally distributed			
	Exclusion criteria: <20 years old congenital heart defects or mechanical reanimation or irradiation			
	Follow-up: 90 days			
	Tewarie et al. (2012), <i>J Cardiothorac Surg</i> [6] Randomized controlled trial, single centre	750 male patients undergoing cardiac surgery with sternotomy	Superficial sternal wound infections	Corset = 8 patients Bandage = 6 patients (P -value not provided)
Group A: Stern-E-Fix corset ($n = 380$)		Deep surgical wound infection	Corset = 4 patients Bandage = 7 patients (P -value not provided)	Patients received support devices for 6 weeks, given from first postoperative day and 96% patients were pleased
Group B: Elastic thorax bandage ($n = 370$)		Sternal dehiscence requiring reoperation	Corset = 1 patient Bandage = 22 patients (P -value not provided)	
Mean follow-up = 8 weeks		Mean ventilation time	Corset = 1.28 days Bandage = 2.5 days ($P = 0.01$)	Similar sternal closure and antibiotic protocol
		Mean hospital stay	Corset = 12.5 days Bandage = 18 days ($P = 0.002$)	
Gorlitzer et al. (2013), <i>Interact CardioVasc Thorac Surg</i> [7] Randomized controlled trial, multicentre	2539 patients, male and female gender, randomized immediately after cardiac surgery to receive Posthorax vest or elastic chest bandage	Superficial wound infection	Vest = 1.55% Bandage = 1.09% ($P = 0.388$)	Intention-to-treat analysis Patients wore the vest for 6 weeks and monitored by specially trained nurse
	Group A: Vest group ($n = 1351$)	Deep sternal complications	Vest = 1.04% Bandage = 2.27% ($P = 0.017$)	Perioperative antibiotic protocol, red cells and plasma transfusion similar for both groups
	Received vest within 48 h = 933 Did not receive in 48 h = 216 Refused vest = 202	Relative risk reduction of suffering a deep sternal complication after vest application	54% lower in vest group	Method of sternal closure not reported
	Group B: Elastic chest bandage ($n = 1176$)			17.8% in treatment group refused to wear the vest

Continued

Table 1: (Continued)

Author, year, journal, study type	Population group	Outcome	Key results	Comments
	No significant risk differences between groups			
	Follow-up: 90 days			
	Exclusion criteria: <20 years old congenital heart defects or mechanical reanimation or irradiation or transplantation			

evaluated whether postoperative an external chest support device reduced the incidence of sternal wound complications in patients following cardiac surgery with sternotomy. Support devices assessed include Posthorax vest, Stern-E-Fix device, Cardibra and the elastic bandage.

Gorlitzer *et al.* [2] evaluated sternal wound complications in adult patients following cardiac surgery by randomizing 455 patients postoperatively to receive Posthorax external support or no support. In this study, 23.2% patients who were randomized to the external support group refused to wear the Posthorax vest due to the close fit and slipping of the vest. These patients were analysed as a separate third group. The patients were followed up for 90 days and the authors report a reduction in overall sternal wound complications (0.6 vs 4.9%; $P = 0.015$) in the vest group. A higher prevalence of diabetes was observed in the Posthorax vest and the refused vest groups compared with the non-vest group. A higher incidence of sternal wound complications (9.4%) was also observed in the refused vest group; however, the incidence of perfusion time exceeding 200 min was higher in this group compared with other groups.

Celik *et al.* [3] reported two studies in their paper. The first study was a retrospective cohort analysis of 842 patients who underwent elective cardiac surgery. The patients were stratified into two groups based on the presence or absence of COPD. Although the technique of sternal closure was similar in all patients, their analysis demonstrated that patients with COPD had higher incidence of sternal dehiscence and deep sternal wound complications (7.9 vs 1.2%, $P < 0.001$). The second study was an RCT with 221 adult patients with COPD who underwent cardiac surgery and were randomized to Posthorax vest postoperatively ($n = 100$) or no vest ($n = 121$). All patients were closed using the Robiscek lateral reinforced sternal closure and were followed up for 6 months postoperatively. They found a significant reduction of overall sternal wound complication (1 vs 11.5%, $P = 0.002$) and deep sternal wound complication (1 vs 9%, $P = 0.02$) in the Posthorax vest group. They analysed and concluded that 15 patients with moderate to severe COPD need to be treated to prevent one sternal wound complication (number needed to treat, NNT = 15).

Naismith and Street [4] performed an RCT involving 20 female patients with bra cup size greater than or equal to C cup and randomly allocated them into Cardibra ($n = 10$) or regular bra ($n = 10$) after cardiac surgery. The aim was to evaluate the effectiveness of Cardibra on sternal wound healing and postoperative pain.

Wound was assessed on 7th, 14th and 42nd day using the Flanagan's wound assessment practical framework by measuring in millimetres, the length of non-approximated wound edges and the area of redness and swelling of the skin around the sternotomy wound. A Likert scale was used for pain scoring on the 1st, 3rd, 5th, 7th, 14th and 42nd day after surgery. They found no significant difference in pain relief or sternal wound complication between groups. However, the sample size in this study was extremely small.

Gorlitzer *et al.* [5] studied the effectiveness of Posthorax vest and the elastic bandage in reducing reoperations due to sternal wound complications. Patients who failed to use the vest were excluded (27.9%) from the final analyses. They observed that the reoperation rates due to sternal wound complications during the 90-day follow-up period were 0.6% in the Posthorax vest group and 3.9% in the elastic bandage group ($P = 0.05$). Total length of hospital stay was also shorter in the Posthorax vest group. They concluded that the need for additional surgical procedures was significantly reduced using the support vest. Unfortunately, some of the patients initially randomized were excluded from the final analysis, but it would have been interesting if this paper had published an analysis on 'intention to treat' basis as well.

Tewarie *et al.* [6] performed an RCT to determine the effect of Stern-E-Fix corset on prevention of sternal wound complications and mediastinitis in patients after cardiac surgery with sternotomy. They randomized 750 male patients to immediately receive either the Stern-E-Fix corset ($n = 380$) or an elastic bandage ($n = 370$) postoperatively. Patients received support devices for 6 weeks, given from first postoperative day, the mean follow-up was 8 weeks and 96% patients were pleased with the design. The sternal closure and antibiotic protocol were similar in all patients and female patients were not included in this study. They observed that only 1 patient in the corset group developed sternal complication requiring a reoperation as opposed to 22 patients in the elastic bandage group. They also found significant reduction in the mean length of hospital stay in the corset group (12.5 vs 18 days; $P = 0.002$). It is important to appreciate that the mean ventilation time was significantly higher in the elastic bandage group (2.5 vs 1.28 days, $P = 0.01$) which may have influenced the results.

Gorlitzer *et al.* [7] reported on a multicentre RCT with 2539 patients assessing the efficacy of Posthorax vest in comparison to elastic bandage in preventing sternal wound complications after cardiac surgery. The Posthorax vest was assigned to 1351 patients

and the elastic bandage to 1176 patients. The Posthorax vest was refused by 17.8% patients after randomization. The patients were followed up for 90 days, and the outcomes were analysed on intention-to-treat basis. They found significant decrease in deep sternal wound complication in the patients randomized to Posthorax group (1.04 vs 2.27%; $P = 0.017$). All complications in the vest group occurred among patients who did not receive or refused the vest.

In these studies, the Stern-E-Fix corset was not evaluated in female patients and a substantial population found the Posthorax vest uncomfortable and therefore refused to wear it.

CLINICAL BOTTOM LINE

Early post-sternotomy use of the available external non-elastic sternal support devices reduces sternal wound complications and may be associated with a shorter length of hospital stay.

Conflict of interest: none declared.

REFERENCES

- [1] Dunning J, Prendergast B, Mackway-Jones K. Towards evidence-based medicine in cardiothoracic surgery: best BETS. *Interact CardioVasc Thorac Surg* 2003;2:405–9.
- [2] Gorlitzer M, Folkmann S, Meinhart J, Posluszny P, Thalmann M, Weiss G *et al.* A newly designed thorax support vest prevents sternum instability after median sternotomy. *Eur J Cardiothorac Surg* 2009;36:335–9.
- [3] Celik S, Kirbas A, Gurer O, Yildiz Y, Isik O. Sternal dehiscence in patients with moderate and severe chronic obstructive pulmonary disease undergoing cardiac surgery: the value of supportive thorax vests. *J Thorac Cardiovasc Surg* 2011;141:1398–402.
- [4] Naismith C, Street A. Introducing the Cardibra: a randomised pilot study of a purpose designed support bra for women having cardiac surgery. *Eur J Cardiovasc Nurs* 2005;4:220–6.
- [5] Gorlitzer M, Wagner F, Pfeiffer S, Folkmann S, Meinhart J, Fischlein T *et al.* A prospective randomized multicenter trial shows improvement of sternum related complications in cardiac surgery with the Posthorax support vest. *Interact CardioVasc Thorac Surg* 2010;10:714–8.
- [6] Tewarie LS, Menon AK, Hatam N, Amerini A, Moza AK, Autschbach R *et al.* Prevention of sternal dehiscence with the sternum external fixation (Stern-E-Fix) corset—randomized trial in 750 patients. *J Cardiothorac Surg* 2012;7:85.
- [7] Gorlitzer M, Wagner F, Pfeiffer S, Folkmann S, Meinhart J, Fischlein T *et al.* Prevention of sternal wound complications after sternotomy: results of a large prospective randomized multicentre trial. *Interact CardioVasc Thorac Surg* 2013;17:515–22.